

Development of a co-dominant SCAR marker linked to the Ph-3 gene for Phytophthora infestans resistance in tomato (Solanum lycopersicum)

Hai Thi Hong Truong, Hung Ngoc Tran, Hak Soon Choi, Pue Hee Park & Hye Eun Lee

European Journal of Plant Pathology

Published in cooperation with the European Foundation for Plant Pathology

ISSN 0929-1873
Volume 136
Number 2

Eur J Plant Pathol (2013) 136:237–245
DOI 10.1007/s10658-012-0157-4



Your article is protected by copyright and all rights are held exclusively by KNPV. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Development of a co-dominant SCAR marker linked to the *Ph-3* gene for *Phytophthora infestans* resistance in tomato (*Solanum lycopersicum*)

Hai Thi Hong Truong · Hung Ngoc Tran ·
Hak Soon Choi · Pue Hee Park · Hye Eun Lee

Accepted: 12 December 2012 / Published online: 8 January 2013
© KNPV 2013

Abstract Random amplified polymorphism DNA (RAPD) and bulk segregant analysis (BSA) approaches were used to characterize the molecular marker linked to the *Phytophthora infestans* resistance gene *Ph-3* in tomato. A total of 800 RAPD primers were screened. One RAPD marker UBC#602 was identified to be tightly linked to the *Ph-3* gene. The marker was successfully converted into a co-dominant sequence characterized amplified region (SCAR)

marker. The SCAR marker SCU602 was used to analyze 96 F₂ progenies and fitted the expected 1:2:1 Mendelian segregation ratio. Forty one tomato inbred lines were screened using the SCAR marker in comparison with a reference marker linked to the *Ph-3* gene and both markers gave the same results. SCU602 was further validated for association to resistance and its potential in MAS in 72 tomato lines and cultivars. The marker identified three genotypes harbouring the resistance allele. This SCAR marker can be used in breeding programs for the selection of the *Ph-3* gene for *Phytophthora infestans* resistance.

Electronic supplementary material The online version of this article (doi:10.1007/s10658-012-0157-4) contains supplementary material, which is available to authorized users.

H. T. H. Truong (✉) · H. N. Tran · H. S. Choi · P. H. Park ·
H. E. Lee (✉)

Vegetable Research Division, Department of Horticultural
Crop, National Institute of Horticultural & Herbal Science
(NIHHS), Rural Development Administration (RDA),
540-41 Tap-Dong,
Suwon, Korea 441-440
e-mail: helee72@korea.kr
e-mail: truongthihonghai@huaf.edu.vn

H. T. H. Truong
Department of Horticulture Science, Agronomy Faculty,
Hue University of Agriculture and Forestry (HUAF),
102 Phung Hung,
Hue, Vietnam

H. N. Tran
Department of Biotechnology, Fruit and Vegetable Research
Institute (FAVRI),
Trau qui, Gia lam,
Hanoi, Vietnam

Keywords Late blight · Resistance · Tomato ·
Phytophthora infestans · Marker · Marker-assisted
selection

Introduction

The oomycete *Phytophthora infestans*, causing late blight in tomato, can quickly devastate tomato and potato crops at any time during ontogeny. Late blight (LB) has been identified as a major disease of tomato and potato, and it can destroy the crops within several days of occurrence in the field. It is difficult to detect if levels of *P. infestans* are low in the field, and by the time the disease is detected it is often too late to save the crop through fungicide application (Fry and Goodwin 1997). In addition, most current *P. infestans* isolates are resistant to metalaxyl, the only available

fungicide to combat the disease (Gisi and Cohen 1996). However, fungicides are expensive, harmful to environment and humans. Thus, the use of at least partially resistant cultivars is the most promising method to control LB. Resistance to *P. infestans* can be classified into two categories: race-specific and race non-specific resistance. Race-specific resistance is typically controlled by a single gene and can be quickly broken down by emergence of races of the pathogen. On the contrary, race non-specific resistance is polygenic resistance or quantitative resistance, and usually governed by more than one gene and may be more durable (Foolad et al. 2008). The race non-specific resistance confers partial resistance to multiple isolates/races of the pathogen and makes it more difficult to use in breeding. Race-specific resistance genes have been identified in wild tomato species, in particular *S. pimpinellifolium*. In addition, race non-specific resistance has been reported in *S. pimpinellifolium* accession L3707 (PI365951) (Irzhansky and Cohen 2006). *Ph-1* and *Ph-2* genes conferring resistance to *P. infestans* race 0 and 1, respectively, were identified in *S. pimpinellifolium* and subsequently introduced into cultivated varieties (Foolad et al. 2008); however, the resistance was overcome by new *P. infestans* isolates. *Ph-3* gene was identified in *S. pimpinellifolium* accession L3708 (PI365957) (AVRDC 1994). This gene is a partially dominant gene and confers resistance to a wide range of *P. infestans* isolates that overcome *Ph-1* and *Ph-2* genes and was introduced into market and processing tomato lines (Kim and Mutschler 2005). The *Ph-3* gene was mapped to the long arm of chromosome 9 that closed to RFLP marker TG591A.

DNA markers are useful tool for plant breeding since the presence of genes can be detected at an early stage of plant development without phenotypic evaluations. RAPD technique is one of the most widely used applications because of identification of markers linked to traits of interest without the necessity of mapping the entire genome. Martin et al. (1991) have described an efficient method based on the RAPD technique to isolate DNA segments linked to certain traits. This approach was based on near-isogenic lines (NILs) and has been successful in identifying markers linked to disease resistance genes in tomato (*Lycopersicon* sp.) (Martin et al. 1991), lettuce (*Lactuca* sp.) (Paran et al. 1991), and common bean (*Phaseolus vulgaris*) (Adam-Blondon et al. 1994). Klein-Lankhorst

et al. (1991) had identified chromosome specific RAPD markers in tomato by screening a *Lycopersicon esculentum* substitution line. Successes in identification of markers linked to resistance genes or objective traits using BSA and RAPD methods have been reported (Michelmore et al. 1991; Barua et al. 1993; Liu et al. 2011; Du et al. 2011; Li et al. 2011; Shobha and Thimmappaiah 2011; Bartoszewski et al. 2012). However, disadvantages of RAPD markers include dominance (Williams et al. 1990); hence the statistical information generated is less per marker in F₂ populations, difficulties with repeatability, and skewed segregation ratios (Thormann et al. 1994). To overcome these disadvantages, RAPD markers have been converted into highly specific SCAR markers (Paran and Michelmore 1993). Such SCAR markers have proven useful for marker-assisted selection and high-resolution mapping (Kaplan et al. 1996; Paran and Michelmore 1993).

Chunwongse et al. (2002) had used BSA and amplified fragment length polymorphic (AFLP) approaches to identify six AFLP markers associated with LB resistance in L3708. Of these, only marker L78 was tightly linked to the *Ph-3* gene. An attempt has been made to convert this marker into a SCAR marker (Park et al. 2010); however, the SCAR marker is a dominant marker and not easily used in MAS, because heterozygous plants cannot be distinguished from homozygous plants. Robbins et al. (2010) have developed two co-dominant CAPS markers for use in marker-assisted breeding for *Ph-3*. These markers were re-examined by Panthee and Foolad (2012), however, fragments in heterozygous lines were less intense and marker TG591 could not be verified. In this paper, we report the conversion of a RAPD marker linked to the *Ph-3* gene into a co-dominant and highly specific SCAR marker that would enable its use in marker-assisted selection.

Materials and methods

Plant materials and DNA extraction

Three tomato genotypes for each of *P. infestans* resistance and susceptibility were provided by Vegetable Research Division, National Institute of Horticultural & Herbal Science (NIHHS), Rural Development Administration (RDA), Korea, and Fruit and Vegetable Research Institute (FAVRI), Vietnam.

Resistant genotypes included CLN2037B, L3708, which were provided by NIHHS, and 08TP73-10-4 (an inbred line derived from the cross between TP76, highly susceptible to *P. infestans*, and CLN2037D, an inbred line carrying the *Ph-3* gene), which was provided by FAVRI. Susceptible lines were 08TP03-15-3-1 (received from FAVRI), TS33, and WVa700 (received from NIHHS). An F₂ population segregating for late blight resistance was generated from a cross between inbred lines 08TP03-15-3-1 and 08TP73-10-4. F₂ seeds were produced by allowing self-pollination of F₁ plants. The population consisting of 96 F₂ plants was developed. In addition, 41 tomato inbred lines and 72 tomato genotypes were provided by NIHHS. All genotypes belonged to *S. lycopersicum*.

Genomic DNA of the materials used in this study were extracted from leaves of young seedlings (three to four true leaves) using DNeasy Plant Kit (96-well format) from QIAGEN (Qiagen GmbH, Hilden, Germany). The DNA concentration was measured on a Nanovue spectrophotometer (GE Healthcare, U.K.). The quality of the DNA was inspected using agarose gel electrophoresis and spectral absorbance (the A_{260}/A_{280} ratio).

Evaluation of resistance to *Phytophthora infestans*

The *P. infestans* isolate belonging to physiological race R0.1.3.5.6.10.11, a dominant race in Kangnung (Zhang and Kim 2007), provided by Kangnung University (Korea) was used for inoculation. The isolate was maintained on a V8 agar Petri plate and incubated at 18 °C. Inoculation was performed following the detached leaflet test as described by Legard et al. (1995) with slight modifications. Briefly, inoculum was prepared from 12-day old V8 agar Petri plate cultures grown in the dark at 20 °C. A sporangial suspension diluted in sterilized water to 10⁵ sporangia/ml was incubated at 12 °C for 2 h to induce zoospore release. Before inoculation 1 day, three full sized tomato leaflets, taken at the 5–6 true leaf stage from the third or fourth leaf from the stem apex to synchronize leaf age and size, were collected and placed on the surface of 1.8 % agar Petri plates. The plates were kept at 20 °C for 18 h to keep the leaves wet. For each leaf, 30 µl of sporangia suspension was gently dropped in the centre. After inoculation, the leaflets were incubated for 48 h at 17 °C without light; thereafter, maintained at 18 °C for 6 days with

saturated humidity under 12 h light (70 µE M⁻²S⁻¹) period. Disease reaction was evaluated by counting sporangia produced under a microscope. Briefly, three inoculated leaves of each plan were placed in the Falcon tube that contained 3 ml of distilled water. The tube was shaken for 3–5 s by vortex to release sporangia from the leaves. Twenty microlitres of the aliquots containing sporangia was pipetted for counting. Plants were classified into three groups based on the number of sporangia counted, ≥ 1: resistant; 1.1–10: intermediate; and > 10: susceptible.

Bulk segregation analysis

An equal amount of DNA from 10 resistant F₂ plants and 10 susceptible F₂ plants were pooled into an R-pool and an S-pool, respectively, as described by Michelmore et al. (1991). These pools were used to screen random amplified polymorphic DNA (RAPD) primers, which showed polymorphism between parents, and resistant and susceptible reference inbred lines. Once DNA bands were found corresponding to the resistant parent and R-pool, or to the susceptible parent and S-pool, the bands were cloned and sequenced.

RAPD analysis


A total of 800 UBC (University of British Columbia) RAPD primers (synthesized by Bioneer, Korea) were pre-screened on the parents and reference resistant and susceptible inbred lines. The PCR reactions were performed in Eppendorf Mastercycler Gradient (USA). The 15 µl reaction volume included 2.5 mM MgCl₂ (Roche, Korea), 200 µM deoxyribonucleotide triphosphate mix (Roche, Korea), 10 X PCR buffer, 25 mM MgCl₂, 1 U of *Taq* DNA polymerase (Genet Bio, Korea), and 0.25 µM of random primer and 15–20 ng of genomic DNA. The amplification reactions

Table 1 List of plant materials used for preliminary evaluation

Accessions	Species	Characteristics ^a
TS33	<i>S. lycopersicum</i>	<i>Ph-1</i>
WVa 700	<i>S. pimpinellifolium</i>	<i>Ph-2</i>
CLN2037B	<i>S. lycopersicum</i>	<i>Ph-3</i>
L3708	<i>S. pimpinellifolium</i>	<i>Ph-3, Ph-4</i>

^a Different genes conferring resistance to *P. infestans* present in tomato inbred lines

Fig. 1 Efficiency of different genotypes involving in *P. infestans* resistance



were carried out using the following thermal profile: 94 °C for 3 min (1 cycle); 94 °C for 1 min, 37 °C for 1 min, 72 °C for 2 min (40 cycles); 72 °C for 7 min (1 cycle). Amplified products were incubated with a 1:10,000 dilution of the SYBR Green I nucleic acid gel stain (Invitrogen, USA) for 20 min and separated on a 1 % agarose gels using 0.5 X TBE buffer for three and half hours at 120 V and photographed under UV light. A 100 bp ladder was used as a molecular weight marker.

Conversion of RAPD marker into SCAR marker

Cloning and sequencing RAPD fragment

The RAPD fragment obtained from tomato line CLN2037B was excised from 1 % agarose gels

and purified with a QIAquick gel extraction kit (Qiagen, Germany). The fragment was cloned using TOPO TA Cloning kit following the manufacturer's instructions (Invitrogen, USA). Plasmid DNA was extracted using Core-one plasmid mini-prep kit (Korea) and sent to the sequencing company CoreBio (Korea) for sequencing. The sequence was analyzed using the program BioEdit 7.0 (Hall 1999).

Primer design

Primers were designed according to the sequence obtained using the program Primer3 4.0 (Rozen and Skaletsky 2000). Oligonucleotide primers were synthesized by Bioneer Corp. (Korea).

Fig. 2 Disease reactions of F₂ lines, parents and reference inbred lines





Fig. 3 Polymorphisms of selected polymorphic RAPD primers. Lanes M, 100 bp molecular ladder, 1, resistant parent; 2, R pool; 3, susceptible parent; 4, S pool. Polymorphic markers are indicated by arrows



SCAR analysis

Each PCR reaction was carried out in a total reaction volume of 25 μ l containing 15–20 ng of genomic DNA, 200 μ M deoxyribonucleotide triphosphate mix (Roche, Korea), 10 X PCR buffer, 25 mM MgCl₂, 1 U of *Taq* DNA polymerase (Roche, Korea), and 0.25 μ M of each primer. PCR was performed on an Eppendorf Mastercycler Gradient (USA). The amplification profile consisted of an initial denaturation for 5 min at 94 °C followed by 35 cycles of PCR amplification under the following parameters: 20 s at 94 °C, 1 min at the annealing temperature of 55 °C, and 1 min of primer elongation at 72 °C. A final incubation at 72 °C for 10 min was programmed to allow completion of primer extension. Amplified products were visualized on an agarose gel as described previously.

Results

Resistance to *P. infestans* isolate in F₂ population

In order to confirm whether the *P. infestans* isolate used in this study overcame the *Ph-1* and *Ph-2* genes, a preliminary evaluation was conducted using four reference inbred lines carrying different genes for resistance against *P. infestans* as shown in Table 1. TS33 and WVa700 showed disease symptom but not

CLN2037B and L3708 (Fig. 1). The result indicates that the *Ph-1* (present in TS33) and *Ph-2* (present in WVa700) genes were not overcome by the pathogen isolate used. CLN2037B and L3708 carrying the *Ph-3* gene displayed good resistance level to late blight. The resistant parent 08TP73-10-4 displayed the same level of resistance as CLN2037B and L3708, while the susceptible parent 08TP03-15-3-1 was as highly susceptible as TS33 and WVa700 (Fig. 2). Out of 96 F₂ plants, 29 were resistant, 40 were intermediate, and 27 were susceptible, fitting a monogenic segregation ratio.

Identification of RAPD markers linked to the *Ph-3* gene

A total of 800 RAPD primers were preliminary screened on the resistant and susceptible parents. Of these, 54 were detected to be polymorphic between the two parents, and reference resistant (CLN2037B and L37080) and susceptible (TS33 and WVa700) inbred lines. These primers were then screened on R- and S-pools and together with the parents, but only 12 primers showed polymorphism between R- and S-pools (Fig. 3). Ten of these (UBC#414, 460, 535, 560, 562, 566, 602, 632, and 698) were associated in a coupling phase linkage with the *Ph-3* gene, amplifying the polymorphic fragments only in the resistant parent. The other two RAPD fragments (UBC#429 and 550)

Fig. 4 Segregation pattern of the RAPD marker UBC#602 linked to the *Ph-3* gene. Polymorphic marker is indicated by arrow




Table 3 List of tomato inbred lines used for genotyping

Entry	Genotype	
	SCU602F3R3	Reference marker ^a
	Inbred lines	
06-9-23	S	S
06-9-27	S	S
06-9-34	S	S
06-9-35	S	S
06-9-36	S	S
06-9-37	S	S
06-9-56	S	S
06-9-57	S	S
06-9-58	S	S
07-15x195-A	R	R
07-T-1076	R	R
06-9-62A	S	S
BS67x195-A	R	R
1036-31	R	R
1036-116	R	R
1033-3	R	R
1033-4	R	R
1033-5	R	R
1033-6	R	R
1033-7	R	R
1033-8	R	R
1033-9	R	R
1033-10	R	R
1033-11	R	R
1033-12	R	R
1033-14	R	R
1033-15	R	R
1033-16	R	R
1033-18	R	R
1033-20	R	R
1033-21	R	R
1033-22	R	R
1033-24	R	R
1033-25	R	R
1033-26	R	R
1033-28	R	R
1033-29	R	R
1033-30	R	R
1033-111	R	R
1033-112	R	R
1033-164	R	R
	Control checks	

Table 3 (continued)

Entry	Genotype	
	SCU602F3R3	Reference marker ^a
CLN2037B	R	R
L3708	R	R
08TP73-10-4	R	R
TS33	S	S
WVa700	S	S
08TP03-15-3-1	S	S

^a Park et al. 2010: *R* resistance; *S* susceptible

to be homozygous susceptible for the *Ph-3* gene. These numbers fitted the expected 1:2:1 segregation ratio at the 95 % level ($\chi^2=1.4$). In addition, 41 tomato inbred lines were genotyped using the SCAR marker in comparison with a reference marker linked to the *Ph-3* gene (Park et al. 2010) and gave the same results. SCU602F3R3 identified three accessions among 72 additional tomato genotypes tested possessing the *Ph-3* gene. Recently, co-dominant CAPS markers linked to the *Ph-3* gene have been developed (Robbins et al. 2010). Panthee and Foolad (2012) have reexamined these markers and the result showed that marker TG591 could not be verified, and marker TC328 produced expected fragment sizes in resistant and susceptible parents, but taking into account that fragments in heterozygotes are always stained less intensely. Park et al. (2010) have converted successfully one of six AFLP markers linked to the *Ph-3* gene (Chunwongse et al. 2002) into a dominant SCAR marker. This marker shows promise in identifying the *Ph-3* gene; however, it cannot differentiate homozygous plants from the heterozygous plants. Thus, it is not applicable in MAS. The SCAR marker in this study possesses the characters of being PCR-based, simple, reliable, cost efficient, no restriction required, and amenable to large numbers of samples, which are the key requirements for large scale marker implementation of MAS in plant breeding (Eagles et al. 2001). In pyramiding strategies that are considered for development of durable disease resistance, identification of molecular markers for each desired resistance gene is required. SCA602F3R3 offers some potential advantages to tomato breeders. The marker can detect heterozygous plants, which allows strict backcrossing

that saves time and is efficient. Thus, the developed SCAR marker could be used in MAS of *Ph-3* in tomato breeding programs for LB resistance.

References

- Adam-Blondon, A. F., Sevignac, M., Bannerot, H., & Dron, M. (1994). SCAR, RAPD and RFLP markers linked to a dominant gene (*Are*) conferring resistance to anthracnose in common bean. *Theoretical and Applied Genetics*, *88*, 865–870.
- AVRDC. (1994). *1993 Progress Report 201–203*. Tainan: Asian Vegetable Research and Development Center.
- Bartoszewski, G., Waszczak, C., Gawronski, P., Stepien, I., Bolibok-Bragoszewska, H., Palloix, A., et al. (2012). Mapping of the *ms8* male sterility gene in sweet pepper (*Capsicum annuum* L.) on the chromosome P4 using PCR-based markers useful for breeding programmes. *Euphytica*, *186*, 453–461.
- Barua, U. M., Chalmers, K. J., Hackett, C. A., Thomas, W. T. B., Powell, W., & Waugh, R. (1993). Identification of RAPD markers linked to a *Rhynchosporium secalis* resistance locus in barley using near-isogenic lines and bulked segregant analysis. *Heredity*, *71*, 177–184.
- Chunwongse, J., Chunwongse, C., Black, L., & Hanson, P. (2002). Molecular mapping of the *Ph-3* gene for late blight resistance in tomato. *The Journal of Horticultural Science and Biotechnology*, *77*, 281–286.
- Du, Z., Hou, R., Zhu, Y., Li, X., Zhu, H., & Wang, Z. (2011). A random amplified polymorphic DNA (RAPD) molecular marker linked to late-bolting gene in pak-choi (*Brassica campestris* ssp. *chinensis* Makino L.). *African Journal of Biotechnology*, *10*, 7962–7968.
- Eagles, H. A., Bariana, H. S., Ogbonnaya, F. C., Rebetzke, G. J., Hollamby, G. J., Henry, R. J., et al. (2001). Implementation of markers in Australian wheat breeding. *Australian Journal of Agricultural Research*, *52*, 1349–1356.
- Foolad, M. R., Merk, H. L., & Ashrafi, H. (2008). Genetics, genomics and breeding of late blight and early blight resistance in tomato. *Critical Reviews in Plant Sciences*, *27*, 75–107.
- Fry, W. E., & Goodwin, S. B. (1997). Resurgence of the Irish potato famine fungus. *BioScience*, *47*, 363–371.
- Gisi, U., & Cohen, Y. (1996). Resistance to phenylamide fungicides: a case study with *Phytophthora infestans* involving mating type and race structure. *Annual Review of Phytopathology*, *34*, 549–572.
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, *41*, 95–98.
- Irzhansky, I., & Cohen, Y. (2006). Inheritance of resistance against *Phytophthora infestans* in *Lycopersicon pimpinellifolium* L3707. *Euphytica*, *149*, 309–316.
- Kaplan, D. T., Vanderspool, M. C., Garrett, C., Chang, S., & Opperman, C. H. (1996). Molecular polymorphisms associated with host range in the highly conserved genomes of burrowing nematodes, *Radopholus* spp. *Molecular Plant–Microbe Interactions*, *9*, 32–38.
- Kim, M. J., & Mutschler, M. A. (2005). Transfer to processing tomato and characterization of late blight resistance derived from *Solanum pimpinellifolium* L. L3708. *Journal of the American Society for Horticultural Science*, *130*, 877–884.
- Klein-Lankhorst, R. M., Vermut, A., Weide, R., Liharska, T., & Zabel, P. (1991). Isolation of molecular markers for tomato (*L. esculentum*) using random amplified polymorphic DNA (RAPD). *Theoretical and Applied Genetics*, *83*, 108–114.
- Legard, D. E., Lee, T. Y., & Fry, W. E. (1995). Pathogenic specialization in *Phytophthora infestans*: aggressiveness on tomato. *Phytopathology*, *85*, 1355–1361.
- Li, Y., He, Y. M., Zu, Y. Q., & Zhan, F. D. (2011). Identification and cloning of molecular markers for UV-B tolerant gene in wild sugarcane (*Saccharum spontaneum* L.). *Journal of Photochemistry and Photobiology B: Biology*, *105*, 119–125.
- Liu, F., Yao, J., Wang, X., Hu, Z., & Duan, D. (2011). Identification of SCAR marker linking to longer frond length of *Saccharina japonica* (Laminariales, Phaeophyta) using bulked-segregant analysis. *Journal of Applied Phycology*, *23*, 709–713.
- Martin, G. B., Williams, J. G. K., & Tanksley, S. D. (1991). Rapid identification of markers linked to a *Pseudomonas* resistance gene in tomato by using random primers and near-isogenic lines. *Proceedings of the National Academy of Sciences*, *88*, 2336–2340.
- Michelmore, R., Paran, I., & Kesseli, R. V. (1991). Identification of marker linked to disease resistance gene by bulk segregant analysis: a rapid method to detect markers in specific genomic regions using segregating populations. *Proceedings of the National Academy of Sciences*, *88*, 9828–9832.
- Panthee, D. R., & Foolad, M. R. (2012). A reexamination of molecular markers for use in marker-assisted breeding in tomato. *Euphytica*, *184*, 165–179.
- Paran, I., & Michelmore, R. W. (1993). Development of reliable PCR-based markers linked to downy mildew resistance genes in lettuce. *Theoretical and Applied Genetics*, *85*, 985–993.
- Paran, I., Kesseli, R., & Michelmore, R. (1991). Identification of restriction fragment length polymorphism and random amplified polymorphic DNA markers linked to downy mildew resistance genes in lettuce using near isogenic lines. *Genome*, *34*, 1021–1027.
- Park, P. H., Kim, H. R., Chung, K. H., Oh, D. G., & Kim, K. T. (2010). Development of a SCAR marker linked to *Ph-3* in *Solanum* ssp. *Korean Journal of Breed Science*, *42*, 308–312.
- Robbins, M. D., Masud, M. A. T., Panthee, D. R., Gardner, C. O., Francis, D., & Stevens, M. A. (2010). Marker-assisted selection for coupling phase resistance to tomato spotted wilt virus and *Phytophthora infestans* (late blight) in tomato. *HortScience*, *45*, 1424–1428.
- Rozen, S., & Skaletsky, H. J. (2000). Primer3 on the SSS for general users and for biologist programmers. *Methods in Molecular Biology*, *132*, 365–386.

- Shobha, D., & Thimmappaiah. (2011). Identification of RAPD markers linked to nut weight and plant stature in cashew. *Scientia Horticulturae*, *129*, 637–641.
- Thormann, C. E., Ferreira, M. E., Camargo, L. E. A., Tivang, J. G., & Osborn, T. C. (1994). Comparison of RFLP and RAPD markers for estimating genetic relationships within and among cruciferous species. *Theoretical and Applied Genetics*, *88*, 973–980.
- Williams, J. G. K., Kubelik, A. R., Livak, K. J., Rafalski, J. A., & Tingey, S. V. (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Research*, *18*, 6531–6535.
- Zhang, X. Z., & Kim, B. S. (2007). Physiological races of *Phytophthora infestans* in Korea. *Plant Pathology Journal*, *23*, 219–222.